

## HRP\* Goat Anti-Rabbit IgG (H+L)

### Description

<b>Catalog Number</b>	BTL0002
<b>Package Size</b>	100ul
<b>Target Name</b>	Rabbit IgG
<b>Reactivity</b>	Rabbit
<b>Purification</b>	The antibody was obtained by affinity purified technique, which removes essentially all goat serum proteins, except the specific antibody for rabbit IgG. Goat anti-rabbit IgG was conjugated to peroxidase by means of a method described by Wilson MB and Nakane PK.
<b>Specificity</b>	The antibody conjugate is specific for rabbit IgG. No cross-reactivity was observed with rabbit IgA, IgM proteins by ELISA.
<b>Immunogen Type</b>	Protein
<b>Immunogen Description</b>	The antiserum was developed in goat using the rabbit IgG as the immunogen
<b>Applications</b>	ELISA WB IHC
<b>Formulation</b>	HRP-labeled Goat IgG in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 50% glycerol.
<b>Storage</b>	Store at -20 °C/1 year
<b>Application</b>	ELISA : 1:5000~1:20000 Western blotting: 1:5000~1:10000 Immunohistochemistry: 1:1000~5000
<b>Background</b>	Wilson MB and Nakane PK. In Immunofluorescence and Related Staining Techniques, Elsevier/North Holland Biomedical Press, Amsterdam, P215 (1978).

### References